

AD-A183 347

OFFICE OF NAVAL RESEARCH

Contract N00014-86-K-0043

R & T Code 413f0001-01

TECHNICAL REPORT No. 44

Theory of Laser-Pulse-Induced Molecular Dynamics: Gas-Phase
Molecular Collisions and Adbond Dynamics

by

Hai-Woong Lee, Sander van Smaalen and Thomas F. George

Prepared for Publication

in

Atomic and Molecular Processes with Short Intense Laser Pulses
edited by A. D. Bandrauk (Plenum, New York), NATO ASI Series

Departments of Chemistry and Physics
State University of New York at Buffalo
Buffalo, New York 14260

July 1987

Reproduction in whole or in part is permitted for any purpose of the
United States Government.

This document has been approved for public release and sale;
its distribution is unlimited.

DTIC
ELECTE
JUL 27 1987
S E D

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/87/TR-44			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York		6b. OFFICE SYMBOL (If applicable)		7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260		7b. ADDRESS (City, State and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Contract N00014-86-K-0043	
8c. ADDRESS (City, State and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		10. SOURCE OF FUNDING NOS.			
		PROGRAM ELEMENT NO.		PROJECT NO.	TASK NO.
					WORK UNIT NO.
11. TITLE Theory of Laser-Pulse-Induced Molecular Dynamics: Gas-Phase Molecular Collisions and Aobond Dynamics					
12. PERSONAL AUTHOR(S) Hai-Woong Lee, Sander van Smaalen and Thomas F. George					
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Yr., Mo., Day) July 1987	
				15. PAGE COUNT 10	
16. SUPPLEMENTARY NOTATION Prepared for publication in <u>Atomic and Molecular Processes with Short Intense Laser Pulses</u> , edited by A. D. Bandrauk (Plenum, New York), NATO ASI Series					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB GR.	MOLECULAR DYNAMICS, MOLECULAR COLLISIONS,		
			LASER INDUCED, ELECTRONIC TRANSITIONS,		
			SHORT PULSES, DESORPTION		
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>A semiclassical study presented here indicates that a sufficiently short and intense pulse can be much more effective in inducing a collisional radiative transition than cw radiation or a long pulse, although the intensity must not be too high because the Rabi oscillation can bring down the probability. For the situation of a molecule physisorbed on a crystalline surface and irradiated by a laser, a master equation approach, used to describe the time evolution of the population of the vibrational adbond levels, shows that for high intensities laser-induced vibrational excitation is the same for pulsed and cw lasers.</p> <p>(Keywords: _____)</p>					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input checked="" type="checkbox"/> DTIC USERS <input type="checkbox"/>			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE NUMBER (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL

Atomic and Molecular Processes with Short Intense Laser Pulses
edited by A. D. Bandrauk (Plenum, New York), NATO ASI Series, in press

THEORY OF LASER-PULSE-INDUCED MOLECULAR DYNAMICS: GAS-PHASE MOLECULAR
COLLISIONS AND ADBOND DYNAMICS

Hai-Woong Lee
Department of Physics
Oakland University
Rochester, Michigan 48063 USA

Sander van Smaalen
Laboratory of Inorganic Chemistry
Materials Science Center
State University of Groningen
Nijenborgh 16
9747 AG Groningen
THE NETHERLANDS

Thomas F. George
Departments of Chemistry and Physics & Astronomy
239 Fronczak Hall
State University of New York at Buffalo
Buffalo, New York 14260 USA

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



INTRODUCTION

A semiclassical study presented here indicates that a sufficiently short and intense pulse can be much more effective in inducing a collisional radiative transition than cw radiation or a long pulse, although the intensity must not be too high because the Rabi oscillation can bring down the probability. For the situation of a molecule physisorbed on a crystalline surface and irradiated by a laser, a master equation approach, used to describe the time evolution of the population of the vibrational adbond levels, shows that for high intensities laser-induced vibrational excitation is the same for pulsed and cw lasers.

GAS-PHASE MOLECULAR COLLISIONS

Most theoretical and experimental work in the past on the subject of molecular collisions occurring in the presence of a laser field has been concerned with the case in which the field consists of continuous wave (cw) radiation. In this case the colliding molecules are influenced by the laser radiation during both the incoming and outgoing trajectories. If, however, the laser field consists of a pulse whose duration is shorter than the collision time (for a typical collision time of 10^{-11} - 10^{-12} s, one must have a (sub)picosecond pulse), only a part of the collision process occurs

in the presence of the laser field. The outcome of the collision therefore may be significantly different depending whether the laser radiation is cw or pulsed. The difference shows up strongly in particular for collision processes involving an electronic transition that occurs via curve crossing, as we have shown in our earlier work.^{1,2} Below we give a brief description of such a transition taking place in the presence of laser radiation, with special attention to the effect of the temporal duration of the laser radiation.³ An experimental study of this effect has recently been reported.

Let us consider an electric-dipole transition induced by collision that occurs in the presence of laser radiation. Within the semiclassical two-state formalism, the Schrödinger equation yields the following coupled equations for the probability amplitudes $c_1(t)$ and $c_2(t)$ for the two states 1 and 2 being considered:

$$i\hbar \frac{dc_1(t)}{dt} = H_{11}c_1 + H_{12}c_2 \quad (1a)$$

$$i\hbar \frac{dc_2(t)}{dt} = H_{21}c_1 + H_{22}c_2 \quad (1b)$$

where H_{11} and H_{22} represent adiabatic potential curves in the dressed-molecule representation for the two states 1 and 2 which, we assume, cross at a certain internuclear distance $R = R_c$ (i.e., $H_{22}(R_c) - H_{11}(R_c) = 0$), the interaction term H_{12} is given by $H_{12} = -\vec{\mu} \cdot \vec{E}_0/2$, $\vec{\mu}$ is the transition moment, and E_0 is the electric field amplitude of the laser radiation. The transition probability $1 \rightarrow 2$ is obtained by solving the coupled equations (1a) and (1b) with the initial conditions $|c_1(0)| = 1$ and $c_2(0) = 0$. These equations, however, cannot be solved analytically in general because the matrix elements H_{ij} vary with time in a complicated manner for a realistic collision system.^{1,3} One therefore must rely on either approximate or numerical solutions. Here we present results of approximate calculations based on the Landau-Zener method along with results of numerical calculations.

Perhaps the simplest approximate solution to Eqs. (1) is provided by the Landau-Zener model. The model assumes that the transition is localized in a narrow neighborhood of the crossing point R_c over which $H_{22} - H_{11}$ varies linearly with R and H_{12} remains constant. The transition probability as the system passes through the crossing point R_c is then given by

$$P = 1 - \exp(-p) \quad (2a)$$

where

$$p = 2\pi[H_{12}(R_c)]^2/\hbar v \gamma [1 - (b/R_c)^2]^{1/2} \quad (2b)$$

b is the impact parameter, γ is the slope of $|H_{22} - H_{11}|$ vs. R at R_c , and v is the relative velocity of the colliding partners at R_c . If the collision occurs in the presence of a short pulse so that the system is illuminated on the way in or out only, the transition probability is given by $P_{\text{in}} = P$. On the other hand, if the system is illuminated both on the way in and out as is the case for cw radiation or a long pulse, the transition probability is given by $P_{\text{cw}} = 2P(1-P)$. To exhibit clearly the intensity dependence of the transition probability, we note that p is proportional to I and let $p = \beta I$.

Then the short-pulse probability and cw probability are given respectively by

$$P_s = 1 - \exp(-\beta I) \quad (3a)$$

$$P_{cw} = 2 \exp(-\beta I)[1 - \exp(-\beta I)] \quad (3b)$$

where, for the case of a short pulse, a square pulse of intensity I is assumed.

In Table 1 we give the Landau-Zener probabilities P_s and P_{cw} for different values of I and b for the case $R_c = 9.5$, $v = 1.83 \times 10^{-4}$, $\gamma = 0.002$ and $\mu = 3$ (all quantities are expressed in atomic units), assuming that the laser beam is linearly polarized in the direction of the relative velocity v of the colliding partners and that the transition moment μ is parallel to the internuclear axis as in a $\Sigma - \Sigma$ transition. The probability P_{cw} as a function of the intensity I exhibits the well-known single-peak structure of the Landau-Zener model, whereas P_s monotonically increases with I . As a result, P_{cw} and P_s differ significantly at high I for which $\beta I \gg 1$. Thus, according to the Landau-Zener model, a short pulse can be much more effective than cw radiation or a long pulse in inducing a transition, if the pulse intensity is sufficiently high.

Table 1. The cw probability P_{cw} and short-pulse probability P_s calculated using the Landau-Zener model.

$b(a.u.)$	$I(W/cm^2)$	10^8	5×10^8	10^9	5×10^9	10^{10}	5×10^{10}	10^{11}	5×10^{11}
1	P_{cw}	0.053	0.23	0.37	0.39	0.13	0.00	0.00	0.00
	P_s	0.027	0.13	0.24	0.74	0.93	1.00	1.00	1.00
5	P_{cw}	0.045	0.20	0.33	0.43	0.18	0.00	0.00	0.00
	P_s	0.023	0.11	0.21	0.69	0.90	1.00	1.00	1.00

In order to check the accuracy of the data presented in Table 1, we also have calculated the transition probability by numerically integrating Eqs. (1). For this calculation, we have assumed that the potential curves satisfy

$$H_{22} - H_{11} = 3 \exp(-0.737R) - 0.0027, \quad (4)$$

which yields $R_c = 9.5$ and $\gamma = 0.002$. We have also assumed a straight-line constant velocity ($= v$) trajectory for simplicity of calculation and have chosen $v = 1.83 \times 10^{-4}$ and $\mu = 3$, as before. The result of our calculation is summarized in Table 2. Comparison of Tables 1 and 2 indicates that the Landau-Zener model yields relatively accurate probabilities at low intensities, but the model fails at high intensities ($I > 10^{11} W/cm^2$). This is mainly due to the inability of the Landau-Zener model to correctly

describe the Rabi oscillation of the molecular system at a high laser intensity.² Since P_{s11} decreases due to the Rabi oscillation as I is increased beyond 10^{11} W/cm², there is a finite range of the intensity over which a short pulse is much more effective than cw radiation. For the model system being considered, this range is 10^{10} - 10^{11} W/cm².

Table 2. The cw probability P_{cw} and short-pulse probability P_s calculated by numerical integration of Eqs. (1).

b(a.u.)	I(W/cm ²)						
		10 ⁸	5×10 ⁸	10 ⁹	5×10 ⁹	10 ¹⁰	5×10 ¹⁰
1	P_{cw}	0.056	0.24	0.39	0.39	0.21	0.15
	P_s	0.029	0.14	0.27	0.73	0.88	0.92
5	P_{cw}	0.034	0.17	0.28	0.41	0.13	0.13
	P_s	0.019	0.091	0.17	0.71	0.93	0.93

b(a.u.)	I(W/cm ²)						
		10 ¹¹	5×10 ¹¹	10 ¹²	5×10 ¹²	10 ¹³	3×10 ¹³
1	P_{cw}	0.24	0.39	0.44	0.48	0.48	0.50
	P_s	0.86	0.73	0.68	0.60	0.61	0.54
5	P_{cw}	0.23	0.40	0.46	0.49	0.48	0.47
	P_s	0.87	0.72	0.64	0.58	0.61	0.63

In view of the recent experimental study³ of the short-pulse effect considered here, we show in Table 3 the total number N_t of transitions per target atom per unit pulse energy, an experimentally observable quantity, for different values of the pulse duration and intensity. Although this quantity was obtained by integrating the Landau-Zener probability over impact parameter, the data shown in the table is expected to be accurate since the Landau-Zener model yields accurate probabilities at the values of I considered here. As can be seen from the table, there is very little difference between long and short pulses at 10^8 - 10^9 W/cm². The difference, however, shows up at 10^{10} W/cm², as a noticeable jump in N_t is seen in going from 10 to 1 ps (the collision time of the system being considered is ~ 2.5 ps).

Table 3. Total number of transitions per target atom per unit pulse energy (in arbitrary units).

$I(W/cm^2)$ \ pulse duration (ps)	1000	100	10	1	0.1
10^8	19.8	19.8	19.9	20.2	20.2
10^9	15.1	15.1	15.7	18.5	18.5
10^{10}	1.45	1.57	2.81	8.75	8.84

In conclusion, the results of our calculations summarized in Tables 1, 2 and 3 indicate that a short pulse can be much more effective in inducing a collisional radiative transition than cw radiation or a long pulse. This short-pulse effect becomes significant if the pulse intensity is sufficiently high that the probability of transition at the crossing point is high (~ 1). On the other hand, the intensity must not be too high because the Rabi oscillation can bring down the probability to a low value at an extremely high intensity.

ADBOND DYNAMICS

Level Populations

We now turn to consider a molecule physisorbed on a crystalline surface and irradiated by a laser. Due to the weak van der Waals bond, the admolecule has a series of vibrational states for the motion perpendicular to the surface. Under certain conditions it is reasonable to assume that a laser, with frequency ω_l , is in resonance with only one pair of these levels, say $|g\rangle$ and $|e\rangle$. The transition frequency is $\omega_{eg} = \omega_e - \omega_g > 0$; the detuning is defined as $\Delta = \omega_l - \omega_{eg}$. Relaxation of the eg adbond occurs due to its interaction with the lattice vibrations of the substrate. In the Markov approximation, this is described through the occurrence of rate constants a_{nk} in a set of first-order differential equations for the level populations and coherences of the reduced density operator of the adbond, the so-called master equation.

For pulsed lasers with a pulse duration Δt short compared to the relaxation times (= the inverse rate constants) of the adbond, the latter can be neglected during the pulse. Then, the time evolution during the pulse is given by the optical Bloch equations,¹⁰

$$\begin{aligned}
 \frac{dR_1(t)}{dt} &= \Delta R_2(t) \\
 \frac{dR_2(t)}{dt} &= -\Delta R_1(t) + \Omega_p(t)R_3(t) \\
 \frac{dR_3(t)}{dt} &= -\Omega_p(t)R_2(t) \quad ,
 \end{aligned}
 \tag{5}$$

where $\Omega(t) = \vec{\mu} \cdot \vec{E}_0(t)/\hbar$ is the time-dependent Rabi frequency; $\vec{E}(t) = \vec{E}_0(t) \cos(\omega_L t)$ is the electric field amplitude of the laser, with slowly-varying envelope $\vec{E}_0(t)$; and $\vec{\mu}$ again is the transition dipole moment. The real-valued quantities R_i are defined by the populations of and coherences between the two adbond levels coupled by the laser, in the rotating frame, according to

$$\begin{aligned} R_1 &= \bar{P}_{ge} + \bar{P}_{eg} \\ R_2 &= -i(\bar{P}_{ge} - \bar{P}_{eg}) \\ R_3 &= P_e - P_g \end{aligned} \quad (6)$$

The total exciting effect up to time t of a laser is characterized by

$$\theta(t) = \int_{-\infty}^t dt' \Omega_p(t') \quad (7)$$

Here, we are only interested in the effect of a complete laser pulse, so that the integral can be taken over a single pulse to give a time-independent parameter θ . Let R_1^0 , R_2^0 and R_3^0 be the values just before the pulse. Then, for zero-detuning, the values right after the pulse are given by

$$\begin{aligned} R_1(\Delta t) &= R_1^0 \\ R_2(\Delta t) &= R_2^0 \cos(\theta) + R_3^0 \sin(\theta) \\ R_3(\Delta t) &= R_3^0 \cos(\theta) - R_2^0 \sin(\theta) \end{aligned} \quad (8)$$

where the start of the pulse is taken as zero point of time, and Δt is the pulse duration. We shall only consider the situation of maximal excitation, which is obtained for $\theta = \pi$, the so-called π -pulses.

Instead of a single pulse, we shall consider a series of equally spaced π -pulses, with interval time t and pulse duration $\Delta t \ll t$. Assume that the system is initially in thermal equilibrium, i.e., $R_1 = 0$, $R_2 = 0$ and $R_3 = R_3(\text{eq})$. Then, after a π -pulse, R_1 and R_2 are zero again, but R_3 is changed into its opposite: $R_3(\Delta t) = -R_3(\text{eq})$. Between two consecutive pulses the adbond evolves in time through its relaxation against the lattice vibrations. This process can be described by a master equation. In the present situation only the populations have to be considered. However, unlike the coupling with the laser, transitions to and from all vibrational states are possible. With the result of the laser pulse as initial condition, the formal solution of the master equation is ($\Delta t \leq t \leq t_p$),

$$\underline{P}(t) = \underline{P}(\text{eq}) + e^{-Wt} \{ \underline{P}(\Delta t) - \underline{P}(\text{eq}) \} \quad (9)$$

where W is the matrix formed by the transition rate constants a_{nk} and has elements

$$W_{nm} = \sum_k a_{nk} \delta_{nm} - a_{mn} \quad (10)$$

$\underline{P}(t)$ is the vector formed by the populations $P_n(t)$, and $\underline{P}(eq)$ refers to the thermal equilibrium distribution, given by $W P_n^{eq} = 0$.

After a number of pulses have passed, the adbond will reach a quasi-steady state, wherein the time evolution of the populations is the same in each interval t_p (see Fig. 1). Then, combining Eqs. (8) and (9), two sets




Fig. 1. Schematic drawing of the time evolution of the level occupations for a pulse sequence in the quasi-steady state. Curve a represents R_3 ; curve b gives $P_g + P_e$; and curve c gives P_n for any other level $n \neq e$ or g .

of equations are obtained, which together determine the constants $P_n(\Delta t)$ in terms of the rate constants a_{nk} and the interval time t_p :

$$\underline{P}(t_p) = \underline{P}(eq) + e^{-Wt_p} \{ \underline{P}(\Delta t) - \underline{P}(eq) \} \quad (11)$$

$$P_n(t_p) = P_n(\Delta t) = P_n^0 \quad (n \neq e, g)$$

$$P_g(t_p) = P_g(\Delta t) = P_g^0 \quad (12)$$

$$P_e(t_p) = P_e(\Delta t) = P_e^0$$

For a continuous wave laser, the Rabi-frequency is a constant, Ω_{cw} . Now, the relaxation processes cannot be neglected during the interaction with the laser. The rate constants a_{nk} give rise to additional terms in the

optical Bloch equations, in a way analogous to their occurrence in the master equation.^{11,12} After some time a true steady state will be reached, in which the populations are determined by

$$\begin{aligned}\sum_k a_{nk} P_n(\infty) &= \sum_k a_{kn} P_k(\infty) \quad (n \neq e, g) \\ \sum_k a_{gk} P_g(\infty) &= \sum_k a_{kg} P_k(\infty) - \frac{1}{2} \frac{\Gamma_{eg}}{\frac{1}{2}\Gamma_{eg}^2 + 2\Delta^2} \{P_g(\infty) - P_e(\infty)\} \\ \sum_k a_{ek} P_e(\infty) &= \sum_k a_{ke} P_k(\infty) + \frac{1}{2} \frac{\Gamma_{eg}}{\frac{1}{2}\Gamma_{eg}^2 + 2\Delta^2} \{P_g(\infty) - P_e(\infty)\},\end{aligned}\quad (13)$$

where $\Gamma_{eg} = \sum_k \{a_{ek} + a_{gk}\}$, and the steady-state values are denoted by $P_n(\infty)$.

Pulsed Laser Versus Continuous Wave Laser

To compare the effect of a pulse train with the effect of a continuous wave (cw) laser on the dynamics of some system, a criterion is needed to compare both lasers. Below two possible criteria will be defined. First, it is required that both lasers have equal average power. Because the Rabi-frequency, $\Omega(t)$, is proportional to the electric field of the laser, the intensity is proportional to $\Omega^2(t)$. Performing the time-average for a series of π -pulses then gives:

$$\Omega_{cw}^2 = \pi^2 / (\Delta t t_p) \quad (14)$$

A second possible criterion is to require the average dissipation from both lasers to be equal. This leads to

$$(P_g^0 - P_e^0)/t_p = (\Omega_{cw}^2 / \Gamma_{eg}) (P_g(\infty) - P_e(\infty)) \quad (15)$$

where P_g^0 , P_e^0 , $P_g(\infty)$ and $P_e(\infty)$ follow from Eqs. (11)-(13).

The efficiency of a pulsed laser over a cw laser to induce some process (e.g., desorption or chemical reaction) will be different for each process. Here we restrict ourselves to the effect on the population of the upper one of the two levels coupled by the laser for a two-level system. A more extensive discussion is given elsewhere.¹¹ Let us define the efficiency ϵ as the ratio of the average excited level population $P(av)$ under pulsed laser irradiation to the steady state population $P_g(\infty)$ under cw laser irradiation. For a two-level system, Eqs. (11)-(13) are easily solved.¹¹ We are interested in the situation where the laser-induced population is substantially larger than the equilibrium thermal occupation of the levels. Then, neglecting the latter, the efficiency is

$$\epsilon = \frac{2}{(1 + e^{-\Gamma t_p})} \frac{(1 - e^{-\Gamma t_p})}{\Gamma t_p} \frac{\frac{1}{2}\Gamma^2 + \Omega_{cw}^2}{\Omega_{cw}^2} \quad (16)$$

For a given integrated intensity of one pulse, the average intensity of the pulse train is proportional to t_p^{-1} . Then a definition for a strong/weak

pulsed laser is $\int_0^{\Delta t} I_p dt \lesssim 1$. Analogously a strong/weak continuous wave laser is defined by (Ω_{cw}^2/Γ^2) being much larger/smaller than one.

Using these definitions, it follows from Eq. (16) that $\epsilon = 1$ in the strong laser limit. This can easily be understood when it is realized that both for the pulsed laser and the cw laser the system is then continuously driven into saturation ($R_3 = 0$). Using criterion 1 (Eq. (14)) leads to

$$\epsilon = \frac{1 - e^{-\Gamma t_p}}{1 + e^{-\Gamma t_p}} \frac{\Gamma^2 + 2\Omega_{cw}^2}{\Gamma^2} \frac{\Gamma \Delta t}{\pi^2} \quad (17)$$

In the weak laser limit this gives $\epsilon = \Gamma \Delta t / \pi^2 \ll 1$, and thus ϵ can be made arbitrarily small by decreasing the pulse duration Δt . The explanation is that the intensity of the pulsed laser is proportional to $\Omega^2 \Delta t$, whereas the exciting effect is only proportional to $\Omega \Delta t$. Decreasing Δt , while $\Omega^2 \Delta t$ is held constant, will decrease $\Omega \Delta t$ too. For a cw laser both the intensity and the exciting effect are proportional to Ω^2 . The latter can be regarded as the limiting form of a pulsed laser with $\Delta t \rightarrow \infty$, and therefore will give the most efficient form of excitation.

For criterion 2, $\epsilon = 1$ is obtained, independent of the laser power. This result reflects the fact that the energy flow into the substrate (is absorbed energy) is proportional to the average occupation of the excited level and not on how P_e varies in time.^{11,13}

ACKNOWLEDGMENTS

This research was supported in part by the National Science Foundation under Grant Nos. CHE-8512406 and CHE-8620274, the Air Force Office of Scientific Research (AFSC), United States Air Force, under Contract No. F49620-86-C-0009, and the Office of Naval Research. The research of S. van Smaalen has been made possible by a fellowship from the Royal Netherlands Academy of Arts and Sciences.

REFERENCES

1. H. W. Lee and T. F. George, J. Phys. Chem. 83:928 (1979).
2. H. W. Lee and T. F. George, Phys. Rev. A, in press.
3. T. Sizer II and M. G. Raymer, Phys. Rev. Lett. 56:123 (1986).
4. H. W. Lee and T. F. George, Phys. Rev. A 29:2509 (1984).
5. S. Efrima, L. Jedrzejek, K. F. Freed, E. Hood, and H. Metiu, J. Chem. Phys. 79:2436 (1983).
6. Z. W. Gortel, H. J. Kreuzer, P. Piercy, and R. Teshima, Phys. Rev. B 27:5066 (1983).
7. B. Fain and S. H. Lin, Surf. Sci. 147:497 (1984).
8. H. F. Arnoldus, S. van Smaalen, and T. F. George, Phys. Rev. B 34:6902 (1986).
9. H. F. Arnoldus, S. van Smaalen and T. F. George, Adv. Chem. Phys., in press.
10. L. Allen and J. H. Eberly, "Optical Resonance and Two-level Atoms," Wiley, New York (1975).
11. S. van Smaalen and T. F. George, Surf. Sci. 183:263 (1987).
12. W. H. Louisell, Quantum Statistical Properties of Radiation Wiley, New York (1973), Ch. 6.
13. S. van Smaalen, H. F. Arnoldus, and T. F. George, Phys. Rev. B 35:1142 (1987).

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. C. B. Harris
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. H. Weaver
Department of Chemical Engineering
and Materials Science
University of Minnesota
Minneapolis, Minnesota 55455

Dr. F. Kutzler
Department of Chemistry
Box 5055
Tennessee Technological University
Cookeville, Tennessee 38501

Dr. A. Reisman
Microelectronics Center of North Carolina
Research Triangle Park, North Carolina
27709

Dr. D. DiLella
Chemistry Department
George Washington University
Washington D.C. 20052

Dr. M. Grunze
Laboratory for Surface Science and
Technology
University of Maine
Orono, Maine 04469

Dr. R. Reeves
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. J. Butler
Naval Research Laboratory
Code 6115
Washington D.C. 20375-5000

Dr. Steven M. George
Stanford University
Department of Chemistry
Stanford, CA 94305

Dr. L. Interante
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Mark Johnson
Yale University
Department of Chemistry
New Haven, CT 06511-8118

Dr. Irvin Heard
Chemistry and Physics Department
Lincoln University
Lincoln University, Pennsylvania 19352

Dr. W. Knauer
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. K.J. Klaubunde
Department of Chemistry
Kansas State University
Manhattan, Kansas 66506

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. Murday
Naval Research Laboratory
Code 6170
Washington, D.C. 20375-5000

Dr. J. B. Hudson
Materials Division
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Theodore E. Madey
Surface Chemistry Section
Department of Commerce
National Bureau of Standards
Washington, D.C. 20234

Dr. J. E. Demuth
IBM Corporation
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. M. G. Lagally
Department of Metallurgical
and Mining Engineering
University of Wisconsin
Madison, Wisconsin 53706

Dr. R. P. Van Duyne
Chemistry Department
Northwestern University
Evanston, Illinois 60637

Dr. J. M. White
Department of Chemistry
University of Texas
Austin, Texas 78712

Dr. D. E. Harrison
Department of Physics
Naval Postgraduate School
Monterey, California 93940

Dr. R. L. Park
Director, Center of Materials
Research
University of Maryland
College Park, Maryland 20742

Dr. W. T. Peria
Electrical Engineering Department
University of Minnesota
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson
Department of Metallurgy and
Materials Science
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. S. Sibener
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Arnold Green
Quantum Surface Dynamics Branch
Code 3817
Naval Weapons Center
China Lake, California 93555

Dr. A. Wold
Department of Chemistry
Brown University
Providence, Rhode Island 02912

Dr. S. L. Bernasek
Department of Chemistry
Princeton University
Princeton, New Jersey 08544

Dr. W. Kohn
Department of Physics
University of California, San Diego
La Jolla, California 92037

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Richard Colton
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Dan Pierce
National Bureau of Standards
Optical Physics Division
Washington, D.C. 20234

Dr. R. Stanley Williams
Department of Chemistry
University of California
Los Angeles, California 90024

Dr. R. P. Messmer
Materials Characterization Lab.
General Electric Company
Schenectady, New York 22217

Dr. Robert Gomer
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Ronald Lee
R301
Naval Surface Weapons Center
White Oak
Silver Spring, Maryland 20910

Dr. Paul Schoen
Code 6190
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. John T. Yates
Department of Chemistry
University of Pittsburgh
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene
Code 5230
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. L. Kesmodel
Department of Physics
Indiana University
Bloomington, Indiana 47403

Dr. K. C. Janda
University of Pittsburgh
Chemistry Building
Pittsburg, PA 15260

Dr. E. A. Irene
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Dr. Adam Heller
Bell Laboratories
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann
Department of Chemistry
University of Southampton
Southampton SO9 5NH
UNITED KINGDOM

Dr. H. Tachikawa
Chemistry Department
Jackson State University
Jackson, Mississippi 39217

Dr. John W. Wilkins
Cornell University
Laboratory of Atomic and
Solid State Physics
Ithaca, New York 14853

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis
Department of Physics
University of California
Irvine, California 92654

Dr. J. T. Keiser
Department of Chemistry
University of Richmond
Richmond, Virginia 23173

Dr. D. Ramaker
Chemistry Department
George Washington University
Washington, D.C. 20052

Dr. R. W. Plummer
Department of Physics
University of Pennsylvania
Philadelphia, Pennsylvania 19104

Dr. J. C. Hemminger
Chemistry Department
University of California
Irvine, California 92717

Dr. E. Yeager
Department of Chemistry
Case Western Reserve University
Cleveland, Ohio 44106

Dr. T. F. George
Chemistry Department
University of Rochester
Rochester, New York 14627

Dr. N. Winograd
Department of Chemistry
Pennsylvania State University
University Park, Pennsylvania 16802

Dr. G. Rubloff
IBM
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. Roald Hoffmann
Department of Chemistry
Cornell University
Ithaca, New York 14853

Dr. Horia Metiu
Chemistry Department
University of California
Santa Barbara, California 93106

Dr. A. Steckl
Department of Electrical and
Systems Engineering
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. W. Goddard
Department of Chemistry and Chemical
Engineering
California Institute of Technology
Pasadena, California 91125

Dr. G.H. Morrison
Department of Chemistry
Cornell University
Ithaca, New York 14853

Dr. P. Hansma
Department of Physics
University of California
Santa Barbara, California 93106

Dr. J. Baldeschwieler
Department of Chemistry and
Chemical Engineering
California Institute of Technology
Pasadena, California 91125